

Forearm support for intensive keyboard users: a field study

Catherine Cook
University of Western Sydney
c.cook@uws.edu.au

Robin Burgess Limerick
University of Queensland
robin@hms.uq.edu.au

Keywords: keyboard, call centre, computer, forearm support

Abstract

The aim of this study was to determine whether providing forearm support when using a normal computer workstation would decrease musculoskeletal discomfort in intensive computer users in a call centre. In a randomised controlled study (n=59), thirty participants were given forearm support for 12 weeks and the remainder were given forearm support for the last 6 weeks of the study. Questionnaires were used at 1, 6 and 12 weeks to obtain information about discomfort, workstation setup, working posture and comfort. At 12 weeks, there were fewer reports of neck, back and wrist discomfort amongst all participants. These findings indicate that for the majority of users, forearm support may be preferable to the "floating" posture implicit in current guidelines for computer workstation setup.

1. Introduction

Working without arm support has been proposed as one of the causal factors of neck and shoulder and arm hand diagnoses (Maeda, 1977; Erdelyi et al., 1988; Hagberg & Sundelin, 1986; Bergqvist, et al., 1995). Despite this, the traditional "floating" posture in which a neutral wrist posture is maintained without supporting the arms is still widely used.

Upper extremity support has been reported to reduce static neck and shoulder muscle load during computer keyboard use (Aaras, et al., 1998; Cook, et al., 2001; Marcus et al., 2002). Aaras et al., (2001) reported a significant decrease in neck, shoulder and back discomfort in a group of computer users who were able to support their whole forearm and hand on a concave workstation. No decrease in discomfort was reported for the distal upper extremity. In a recent prospective epidemiological study of computer users, Marcus et al., (2002) reported use of the keyboard placed more than 12cm from the edge of the desk was associated with a lower risk of hand arm symptoms. In a recent laboratory study, forearm support using a conventional desk was also found to result in significantly less ulnar deviation, less time spent in an extreme wrist posture and fewer reports of discomfort (Cook et al., 2001).

Supporting the forearm on the work surface may decrease discomfort, decrease muscular load of the neck and shoulders and decrease harmful wrist postures, thereby creating a beneficial posture for keyboard and mouse users. However, the benefits of providing arm support during keyboard and mouse use in a conventional workstation previously have been described only in the laboratory setting (Cook et al., 2001).

As the provision of specialised equipment such as concave desks is not always practical owing to cost, the effect of adjusting a conventional workstation to allow forearm support during keyboard operation requires evaluation in a field setting. The aim of this study was to determine whether adjusting a conventional workstation to enable forearm support during computer use decreases reports of neck/shoulder or wrist/hand musculoskeletal discomfort in intensive computer users in a field setting.

2. Methods

Participants were experienced keyboard users who worked in a newspaper call centre for at least 15 hours per week. Anyone receiving treatment for musculoskeletal discomfort was excluded from the study. There were 59 volunteers (54 female, 5 male). The average age was 39 years (range 21-68years). Average duration of computer usage was 29 hours per week (SD 6.48 hours). Mean duration of typing experience was 15 years (SD 11.9). Call centre staff are responsible for keying all information for classified advertisements. Information is taken either via the telephone or via email. The mouse is used frequently by those receiving advertisements via email, but rarely by those taking telephone calls. Management reported that about 75% of work time is spent keying. Computer workstations consist of adjustable-height desks, chairs and monitors and telephone headsets. Employees do not necessarily use the same workstation each day.

2.1. Design and measures

The study was a randomised controlled field trial lasting 12 weeks. Participants were randomised to two groups: the forearm support Group 1 (intervention at week 1, n=30) and Group 2 (intervention at week 6, n=29).

Procedures

Week 1: A six-page self-report questionnaire was completed by each participant. The first section requested information on the participant's work patterns, including hours of computer and computer mouse use at work and at home, break frequency and duration, and exercise (Cook et al., 2000). The second part of the questionnaire was based on the Nordic Questionnaire (Kuorinka et al., 1987). Participants were asked to record whether they had experienced musculoskeletal trouble (ache pain or discomfort) in the neck, shoulder, wrist/hand, forearm or back either in the past 12 months or within the last 7 days. Individual workstation assessments were conducted on each participant. Both groups received education about workstation set up and working posture. All workstations were adjusted according to Australian Standard AS3590.2 (Standards Australia, 1990). Measurements of workstation dimensions were recorded before and after adjusting workstations. Wrist rests, if used, were left in situ after workstation setup (31% of participants).

Group 1: Following questionnaire administration at week 1, workstations were adjusted to enable each Group 1 participant (n=30) to support their forearms (but not elbows) on the desk surface, maintaining neutral shoulder elevation. The keyboard was positioned so that the top row of keys was level with fingertips when the forearms were supported comfortably on the worksurface. The positions of the keyboard, desk and chair heights were recorded and marked with tape. The mouse was positioned next to the keyboard, so that at least half of the forearm was supported on the desk while working. Participants were monitored for the first few hours after the changes to their working posture to ensure that they were not adopting postures of trunk flexion, shoulder elevation or increased wrist extension.

Group 2: Where required, adjustments to desk, chair and monitor height were made according to Australian Standards (Standards Australia, 1990). The keyboard and mouse position was recorded.

Week 6: Discomfort questionnaires were again completed by both groups and the forearm support posture (as described above) was introduced to the people in Group 2.

Participants were monitored frequently after introducing the forearm support posture and weekly thereafter to ensure the consistency of working postures. Participants were requested to report any increases in discomfort.

Week 12: All participants completed the musculoskeletal symptom questionnaire and answered additional generated questions about the forearm support posture. A goniometer was used to measure shoulder flexion when the hands were positioned on the keyboard, with the fingertips touching the most distant row of keys (numeric or function keys) and the position of the keyboard remeasured.

2.2. Data Analysis

Symptom prevalence was calculated before intervention, at 6 weeks and 12 weeks after intervention. Differences within the groups at each of the measurement occasions was analysed using McNemar chi-square (Norman & Streiner, 1994). Comparisons were made between the two groups at each measurement occasion.

3. Results

Within a week of intervention, nine participants (15%) withdrew from using forearm support either due to discomfort (4), or difficulty maintaining the posture (4). The forearm support posture was discontinued by the experimenter for one participant who was observed to adopt a posture of increased trunk flexion, due to her abdominal depth. The discomfort reported by these participants on discontinuation of the forearm support posture have been included in the analysis. Two other participants were eliminated from the study, due to unexpected absences. Of the 48 participants who continued, 23 completed 12 weeks and 25 participants completed 6 weeks with forearm support.

3.1. Musculoskeletal discomfort

Before intervention (n=57)

All but one participant reported having had musculoskeletal discomfort in one or more body region in the past 12 months, with 75% reporting discomfort in the 7 days preceding study commencement. There were no differences in overall symptom reporting between the 2 groups prior to intervention (chi 0.478, $p = 0.49$). One third of the participants reported having received treatment for their discomfort within the past 12 months.

Week 6

Group 1 - forearm support group

The proportion of participants reporting discomfort in one or more body area in the previous 7 days decreased from 79% in week 1 to 62% in week 6 (chi = 0.468, $p = 0.227$).

Group 2 - no forearm support

Reports of discomfort increased slightly for all body regions for this group of participants, with overall discomfort increasing from 71% in week 1 to 75% in week 6.

Week 12 (n=57)

There was a significant decrease in overall discomfort for the whole group (ie discomfort in any body region) between week 1 (75%) and 12 (45%) (chi = 0.773, $p = 0.002$). The proportion of participants reporting neck (chi = 5.05, $p = 0.008$), wrist (chi = 0.93, $p = 0.021$) and forearm (chi = 0.062, $p = 0.049$) discomfort decreased between weeks 1 and 12. Shoulder and back discomfort had also decreased although these changes were not statistically significant (shoulder chi = 2.9, $p = 0.36$), (back chi = 4.4, $p = 0.61$). There were no significant differences in reporting of discomfort between the 2 groups (chi 0.139, $p = 0.68$) (Figure 1).




Figure 1: Proportion of participants reporting symptoms per body region.

3.2. Subjective reports:

Of the group (n=48) who completed the study, 2 participants reported that they had used the forearm support posture minimally during the study. They reported that they had not found the position comfortable, and found that because they changed workstation between shifts it was too difficult to set up. Thirty two percent of participants reported using forearm support some of the time, and 64% used forearm support all of the time. The main reason given for not always using the forearm support posture was speed and difficulty getting used to the new technique.

Two participants reported that they found the forearm support position less comfortable, 18% reported the same level of comfort and 72% found it more comfortable than their previous working posture. Reasons given for increased comfort included being able to rest the arms resulting in less pain (3), less strain (4), a more balanced posture (2) and more space (1). Three people reported adjustments other than forearm support such as their leg position as a result of altered chair height, or monitor height as being reasons for their increased comfort. Difficulty adjusting to the new posture was reported by 4 people.

3.3. Mouse use

Mouse users fell into 2 groups with 54% of participants using their mouse most of the day, with the remainder using their mouse less than one hour per day. There were no significant differences in discomfort for any body region before or after intervention between these 2 groups.

3.4. Wrist rests

Fifteen participants (26%) used solid foam wrist rests prior to commencement of the study. On completion of the study, this had increased to 26 participants (46%). The researcher introduced the use of wrist rests to 6 participants during the initial set up into forearm support posture. These participants were observed to rest their wrists on the worksurface while keying resulting in wrist extension. The other 5 participants commenced using wrist rests due to reports of pressure on the forearm from the edge of the desk. The majority used the wrist rests adjacent to the keyboard, with 6 using the wrist rests under their forearm close to the edge of the desk.

3.5. Position of the keyboard

The mean angle of shoulder flexion when working in the forearm support posture was 21 degrees (SD 9.3, range 5-50 degrees), with 4 subjects positioned in more than 30 degrees of shoulder flexion.

The mean distance of the keyboard from the desk edge increased significantly ($t = -9.05$, $p = 0.00$, $df = 47$) from 95mm (SD 50.46) prior to set up, to 168mm (SD 50.7) at 12 weeks. The computer mouse was positioned adjacent to the keyboard enabling forearm support during mouse use.

4. Discussion

There was a significant decrease in the frequency of reported neck, back, forearm and wrist discomfort following use of forearm support when using a conventional desk. A reduction in reported discomfort occurred within 6 weeks of using forearm support for both groups. Nearly two thirds of participants reported using the forearm support posture all of the time, with about three-quarters reporting it more comfortable than their previous work posture. While 20% of participants reverted to a floating posture due to discomfort or dislike of the forearm support position, the majority of users reported an improvement in comfort.

The findings of fewer reports of neck, shoulder and back discomfort are consistent with those of a previous study on forearm support, with some differences in symptom reporting noted between the two studies for the distal upper extremity (Aaras et al., 1998; Aaras, et al., 2001).

Significant reductions in reported wrist and forearm discomfort at 12 weeks contrasts with the lack of difference in discomfort intensity after initial intervention in a previous study (Aaras et al., 1998), with a tendency for increased forearm hand pain over 6 years reported for two of the study groups (Aaras et al., 2001). In the current study, one third of participants used wrist rests on study commencement, with a further quarter of the subjects provided with wrist supports during the study either because of observed wrist extension or reported discomfort from the desk edge. Although not measured, regular observation was conducted by the researcher to check that participants were not working in wrist extension. Due to the possible relationship between wrist rests and increases in carpal tunnel pressure (Horie, et al., 1993), wrist supports were placed under the forearm rather than adjacent to the keyboard where possible (Hedge, 2001). Consistent with a previous laboratory study (Cook et al., 2001) whilst working with forearm support, participants were observed to pivot their forearms rather than anchoring their wrists and working in ulnar deviation. Although Marcus et al., (2002) reported an association between use of wrist rests and hand arm symptoms and disorders, this was for the group who used their keyboard less than 12 cm from the edge of the desk, a working position which they associated with increased risk. The current study confirms those of Marcus et al., (2002), wrist rest use and forearm support were not related to an increase in wrist discomfort.

Most participants in this study were working in less than 30 degrees of shoulder flexion. Working in shoulder flexion of more than above 30 degrees has been associated with increased shoulder discomfort or decreased performance in the unsupported arm (Sauter et al., 1991; Straker, et al., 1997). However, the effect of this shoulder posture in the supported arm has not been reported. Marcus et al., (2002) reported an association between inner elbow angles of >121 degrees and a decreased risk of neck shoulder symptoms. Although not examined separately, this elbow posture was associated with a mean shoulder flexion of 38 degrees.

There was a non-significant increase in discomfort reported at 6 weeks for Group 2 (prior to intervention). Both groups had received education regarding correct workstation setup at the commencement of the study. As most workstations were well adjusted, minimal changes were made to the workstations of the control group. The positive benefit of education alone has been questioned in some studies. An increase in discomfort was reported post intervention in a study where self education had been a major component (Coury, 1998). An increase in forearm/hand pain was reported following a traditional education programme (Bohr, 2000). Lewis et al., (2001) reported that education by professionals with subsequent workstation adjustment by employees did not result in a significant decrease in musculoskeletal discomfort.

5. Conclusion

The results of this study confirm that use of forearm support has a number of advantages over a traditional floating posture and should be considered as an alternate working posture for keyboard users.

6. References

- Aaras, A., Horgen, G., Bjorset, H.-H., Ro, O., & Thoresen, M. (1998). Musculoskeletal, visual and psychosocial stress in VDU operators before and after multidisciplinary ergonomic interventions. *Applied Ergonomics*, 29(5), 335-360.
- Aaras, A., Horgen, G., Bjorset, H.-H., Ro, O., & Walsoe, H. (2001). Musculoskeletal, visual and psychosocial stress in VDU operators before and after multidisciplinary ergonomic interventions. A 6 years prospective study. *Applied Ergonomics*, 32, 559-571.
- Bergqvist, U., Wolgast, E., Nilsson, B., & Voss, M. (1995). Musculoskeletal disorders among visual display terminal workers: individual, ergonomic and work organisational factors. *Ergonomics*, 38(4), 763-776.
- Bohr, P. (2000). Efficacy of Office Ergonomics Education. Paper presented at the IEA 2000/HFES 2000 Congress, San Diego.
- Cook, C., Burgess Limerick, R., & Papalia, S. (2001). The effect of forearm support on posture and muscle activity of the neck and upper extremity during computer keyboard use. Unpublished manuscript.
- Cook, C., Burgess-Limerick, R., & Chang, S. (2000). The prevalence of neck and upper extremity musculoskeletal symptoms in computer mouse users. *International Journal of Industrial Ergonomics*, 26(3), 347-356.
- Coury, H. (1998). Self-administered preventive programme for sedentary workers: reducing musculoskeletal symptoms or increasing awareness. *Applied Ergonomics*, 29(6), 415-421.
- Erdelyi, A., Sihvonen, T., Helin, P., & Hanninen, O. (1988). Shoulder strain in keyboard workers and its alleviation by arm supports. *International Archives of Occupational and Environmental Health*, 60, 119-124.
- Hagberg, M., & Sundelin, G. (1986). Discomfort and load on the upper trapezius muscle when operating a word processor. *Ergonomics*, 29(12), 1637-1645.
- Hedge, A. (2001). Ergonomic Guidelines for arranging a Computer Workstation - 10 steps for users [Online]. Available: <http://ergo.human.cornell.edu/ergoguide.html> [2001, September 30].
- Horie, S., Hargens, A., & Rempel, D. (1993). The effect of keyboard wrist rest in preventing carpal tunnel syndrome. Paper presented at the Proceedings of American Public Health Association Annual meeting, San Francisco.
- Kuorinka, I., Jonsson, B., Kilbom, A., H Vinterberg, F Biering-Sorensen, G Anderson, & Jorgensen, K. (1987). Standardised Nordic Questionnaires for the analysis of musculoskeletal symptoms. *Applied Ergonomics*, 18(3), 233-237.
- Lewis, R., Fogelman, M., Deeb, J., E Crandell, E., & Agopswicz, D. (2001). Effectiveness of a VDT training program. *International Journal of Industrial Ergonomics*, 27, 119-131.
- Maeda, K. (1977). Occupational cervicobrachial disorder and its causative factors. *Journal Human Ergology*, 6, 193-202.
- Marcus, M., Gerr, F., Monteilh, C., Ortiz, D., Gentry, E., Cohen, S., Edwards, A., Ensor, C., & Kleinbaum, D. (2002). A Prospective Study of Computer Users: II Postural Risk Factors for Musculoskeletal Symptoms and Disorders. *American Journal of Industrial Medicine*, 41, 236-249.

- Norman, G., & Streiner, D. (1994). *Biostatistics. The Bare Essentials*. St Louis: Mosby Year Book.
- Sauter, S., Schleifer, L., & Knutsen, S. (1991). Work posture, workstation design, and musculoskeletal discomfort in a VDT data entry task. *Human Factors*, 33(2), 151-167.
- Standards Australia. (1990). *Screen-based workstations. Part 2: Workstation furniture (AS3590.2-1990)*: Standards Australia.
- Straker, L. M., Pollock, C., & Mangharam, J. E. (1997). The effect of shoulder posture on performance, discomfort and muscle fatigue whilst working on a visual display unit. *International Journal of Industrial Ergonomics*, 20, 1-10.