

# Directed Scale-Free Graphs

Béla Bollobás\*

Christian Borgs<sup>†</sup>

Jennifer Chayes<sup>‡</sup>

Oliver Riordan<sup>§</sup>

## Abstract

We introduce a model for directed scale-free graphs that grow with preferential attachment depending in a natural way on the in- and out-degrees. We show that the resulting in- and out-degree distributions are power laws with different exponents, reproducing observed properties of the world-wide web. We also derive exponents for the distribution of in- (out-) degrees among vertices with fixed out- (in-) degree. We conclude by suggesting a corresponding model with hidden variables.

## 1 Introduction

Recently many new random graph models have been introduced and analyzed, inspired by certain common features observed in many large-scale real-world graphs such as the ‘web graph’, whose vertices are web pages (or sites), with a directed edge for each link between two web pages. For an overview see the survey papers [2] and [15]. Other graphs studied are the ‘internet graph’ [18], movie actor [28] and scientific [25] collaboration graphs, cellular networks [21] and many other examples.

In addition to the ‘small-world phenomenon’ of logarithmic diameter investigated originally by Strogatz and Watts [28], one of the main observations is that the graphs are ‘scale-free’ (see [5, 7, 24] and the references therein); the distribution of vertex degrees follows a power law, rather than the Poisson distribution of the classical random graph models  $G(n, p)$  and  $G(n, M)$  [16, 17, 19], see also [9].

Many models have been suggested to explain this and other features of the graphs studied. One of the basic ideas is the combination of growth with ‘preferential attachment’; the graph grows one vertex

at a time, and edges are added, perhaps only from the new vertex to old vertices, or perhaps also between old vertices, where the old vertices involved are chosen with probabilities proportional to their degrees. One of the simplest and earliest models is that outlined by Barabási and Albert in [5], made precise in [11]. The degree sequence of this model was analyzed heuristically in [5, 6], and rigorously in [12]. Many generalizations have been suggested and studied heuristically; a few have been analyzed precisely, see [27]. In a complicated paper Cooper and Frieze [14] have analyzed rigorously a very general version of the model allowing for (finite) distributions of out-degrees and mixtures of uniform and preferential attachment.

The models mentioned above essentially describe undirected graphs. The only exception is [14], where the authors treat either in-degrees or out-degrees, but not both simultaneously; a full treatment of directed graphs was announced there, but has not yet appeared. However, in many contexts – for example the web graph – it is natural to look at directed graphs, and to study the (often different) power laws for in- and out-degrees. Here we propose a very natural model of directed web graphs and show that it gives power laws consistent with those that have been observed in the world-wide web.

Before turning to our model let us briefly mention two rather different kinds of model: Newman, Strogatz and Watts [26], Aiello, Chung and Lu [1] and other groups have studied random graphs chosen by first fixing the (scale-free) degree distribution, and then choosing a graph with this degree distribution. This is very different from our aim here, which is to *explain* the power-law distributions. Also, instead of preferential attachment, copying models have been studied [22, 24]; for the web this is very natural, and something like this is needed to explain the high density of small subgraphs. Again, however, such models are not what we are concerned with here: firstly they do not model out-degree distributions (the out-degrees are fixed). Secondly, they are rather specific. By keeping the model simple we hope that it can give insight into many different scale-free graphs, rather than just the web graph. Also, Cooper and Frieze [14] note that for (in-)degree distribution there is little difference between copying and preferential attachment.

\*Department of Mathematical Sciences, University of Memphis, Memphis TN 38152 and Trinity College, Cambridge CB2 1TQ, UK. Research supported by NSF grant DSM 9971788 and DARPA grant F33615-01-C-1900, and undertaken during a long term visit to Microsoft Research. [bollobas@msci.memphis.edu](mailto:bollobas@msci.memphis.edu)

<sup>†</sup>Microsoft Research, One Microsoft Way, Redmond, WA 98122. [borgs@microsoft.com](mailto:borgs@microsoft.com)

<sup>‡</sup>Microsoft Research, One Microsoft Way, Redmond, WA 98122. [jchayes@microsoft.com](mailto:jchayes@microsoft.com)

<sup>§</sup>Trinity College, Cambridge CB2 1TQ, UK. Research undertaken while visiting Microsoft Research. [O.M.Riordan@dpms.cam.ac.uk](mailto:O.M.Riordan@dpms.cam.ac.uk)

## 2 The model

We consider a directed graph which grows by adding single edges at discrete time steps. At each such step a vertex may or may not also be added. For simplicity we allow multiple edges and loops. More precisely, let  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta_{in}$  and  $\delta_{out}$  be non-negative real numbers, with  $\alpha + \beta + \gamma = 1$ . Let  $G_0$  be any fixed initial directed graph, for example a single vertex without edges, and let  $t_0$  be the number of edges of  $G_0$ . (Depending on the parameters, we may have to assume  $t_0 \geq 1$  for the first few steps of our process to make sense.) We set  $G(t_0) = G_0$ , so at time  $t$  the graph  $G(t)$  has exactly  $t$  edges, and a random number  $n(t)$  of vertices. In what follows, to *choose* a vertex  $v$  of  $G(t)$  according to  $d_{out} + \delta_{out}$  means to choose  $v$  so that  $\Pr(v = v_i)$  is proportional to  $d_{out}(v_i) + \delta_{out}$ , i.e., so that  $\Pr(v = v_i) = (d_{out}(v_i) + \delta_{out}) / (t + \delta_{out}n(t))$ . To *choose*  $v$  according to  $d_{in} + \delta_{in}$  means to choose  $v$  so that  $\Pr(v = v_i) = (d_{in}(v_i) + \delta_{in}) / (t + \delta_{in}n(t))$ . Here  $d_{out}(v_i)$  and  $d_{in}(v_i)$  are the out-degree and in-degree of  $v_i$ , measured in the graph  $G(t)$ .

For  $t \geq t_0$  we form  $G(t+1)$  from  $G(t)$  according to the following rules:

(A) With probability  $\alpha$ , add a new vertex  $v$  together with an edge from  $v$  to an existing vertex  $w$ , where  $w$  is chosen according to  $d_{in} + \delta_{in}$ .

(B) With probability  $\beta$ , add an edge from an existing vertex  $v$  to an existing vertex  $w$ , where  $v$  and  $w$  are chosen independently,  $v$  according to  $d_{out} + \delta_{out}$ , and  $w$  according to  $d_{in} + \delta_{in}$ .

(C) With probability  $\gamma$ , add a new vertex  $w$  and an edge from an existing vertex  $v$  to  $w$ , where  $v$  is chosen according to  $d_{out} + \delta_{out}$ .

The probabilities  $\alpha$ ,  $\beta$  and  $\gamma$  clearly should add up to one. To avoid trivialities, we will also assume that  $\alpha + \gamma > 0$ . When considering the web graph we take  $\delta_{out} = 0$ ; the motivation is that vertices added under step (C) correspond to web pages which purely provide content - such pages never change, are born without out-links and remain without out-links. Vertices added under step (A) correspond to usual pages, to which links may be added later. While mathematically it seems natural to take  $\delta_{in} = 0$  in addition to  $\delta_{out} = 0$ , this gives a model in which every page not in  $G_0$  has either no in-links or no out-links, which is rather unrealistic and uninteresting! A non-zero value of  $\delta_{in}$  corresponds to insisting that a page is not considered part of the web until something points to it, typically one of the big search engines. It is natural to consider these edges from search engines separately from the rest of the graph, as they are of a rather different nature; for the same reason it is natural not to insist that  $\delta_{in}$  be an integer. We include the parameter  $\delta_{out}$  to

make the model symmetric with respect to reversing the directions of edges (swapping  $\alpha$  with  $\gamma$  and  $\delta_{in}$  with  $\delta_{out}$ ), and because we expect the model to be applicable in contexts other than that of the web graph.

Our model allows loops and multiple edges; there seems no reason to exclude them. However, there will not be very many, so excluding them would not significantly affect our conclusions.

Note also that our model includes (a precise version of) the  $m = 1$  case of the original model of Barabási and Albert as a special case, taking  $\beta = \gamma = \delta_{out} = 0$  and  $\alpha = \delta_{in} = 1$ . We could introduce more parameters, adding  $m$  edges for each new vertex, or (as in [14]) a random number with a certain distribution, but one of our aims is to keep the model simple, and the main effect, of varying the overall average degree, can be achieved by varying  $\beta$ .

## 3 Analysis

Having decided on the model it is not hard to find the power laws for in- and out-degrees. Throughout we fix constants  $\alpha, \beta, \gamma \geq 0$  summing to 1 and  $\delta_{in}, \delta_{out} \geq 0$ , and set

$$c_1 = \frac{\alpha + \beta}{1 + \delta_{in}(\alpha + \gamma)} \quad \text{and} \quad c_2 = \frac{\beta + \gamma}{1 + \delta_{out}(\alpha + \gamma)}.$$

We also fix a positive integer  $t_0$  and an initial graph  $G(t_0)$  with  $t_0$  edges. Let us write  $x_i(t)$  for the number of vertices of  $G(t)$  with in-degree  $i$ , and  $y_i(t)$  for the number with out-degree  $i$ .

Note that the in-degree distribution becomes trivial if  $\alpha\delta_{in} + \gamma = 0$  (all vertices not in  $G_0$  will have in-degree zero) or if  $\gamma = 1$  (all vertices not in  $G_0$  will have in-degree 1), while for  $\gamma\delta_{out} + \alpha = 0$  or  $\alpha = 1$  the out-degree distribution becomes trivial. We will therefore exclude these cases in the following theorem.

**THEOREM 3.1.** *Let  $i \geq 0$  be fixed. There are constants  $p_i$  and  $q_i$  such that  $x_i(t) = p_i t + o(t)$  and  $y_i(t) = q_i t + o(t)$  hold with probability 1. Furthermore, if  $\alpha\delta_{in} + \gamma > 0$  and  $\gamma < 1$ , then as  $i \rightarrow \infty$  we have*

$$p_i \sim C_{IN} i^{-X_{IN}},$$

where  $X_{IN} = 1 + 1/c_1$  and  $C_{IN}$  is a positive constant. If  $\gamma\delta_{out} + \alpha > 0$  and  $\alpha < 1$ , then as  $i \rightarrow \infty$  we have

$$q_i \sim C_{OUT} i^{-X_{OUT}},$$

where  $X_{OUT} = 1 + 1/c_2$  and  $C_{OUT}$  is a positive constant.

In the statement above, the  $o(t)$  notation refers to  $t \rightarrow \infty$  with  $i$  fixed, while  $a(i) \sim b(i)$  means  $a(i)/b(i) \rightarrow 1$  as  $i \rightarrow \infty$ .

*Proof.* Note first that if the initial graph has  $n_0$  vertices then  $n(t)$  is equal to  $n_0$  plus a Binomial distribution with mean  $(\alpha + \gamma)(t - t_0)$ . It follows from standard results (e.g., the Chernoff bound) that there is a positive constant  $c$  such that for all sufficiently large  $t$  we have

$$(3.1) \quad \Pr\left(|n(t) - (\alpha + \gamma)t| \geq t^{1/2} \log t\right) \leq e^{-c(\log t)^2}.$$

In particular, the probability above is  $o(t^{-1})$  as  $t \rightarrow \infty$ .

We consider how the vector  $(x_0(t), x_1(t), \dots)$ , giving for each  $i$  the number of vertices of in-degree  $i$  in the graph  $G(t)$ , changes as  $t$  increases by 1. Let  $G(t)$  be given. Then with probability  $\alpha$  a new vertex with in-degree 0 is created at the next step, and with probability  $\gamma$  a new vertex with in-degree 1 is created. More importantly, with probability  $\alpha + \beta$  the in-degree of an old vertex is increased. In going from  $G(t)$  to  $G(t+1)$ , from the preferential attachment rule, given that we perform operation (A) or (B), the probability that a particular vertex of in-degree  $i$  has its in-degree increased is exactly  $(i + \delta_{in})/(t + \delta_{in}n(t))$ . Since the chance that we perform (A) or (B) is  $\alpha + \beta$ , and since  $G(t)$  has exactly  $x_i(t)$  vertices of in-degree  $i$ , the chance that one of these becomes a vertex of in-degree  $i+1$  in  $G(t+1)$  is exactly

$$(\alpha + \beta)x_i(t) \frac{i + \delta_{in}}{t + \delta_{in}n(t)},$$

so with this probability the number of vertices of in-degree  $i$  decreases by 1. However, with probability

$$(\alpha + \beta)x_{i-1}(t) \frac{i - 1 + \delta_{in}}{t + \delta_{in}n(t)}$$

a vertex of in-degree  $i - 1$  in  $G(t)$  becomes a vertex of in-degree  $i$  in  $G(t)$ , increasing the number of vertices of in-degree  $i$  by 1. Putting these effects together,

$$(3.2) \quad \mathbf{E}\left(x_i(t+1) \middle| G(t)\right) = x_i(t) + \frac{\alpha + \beta}{t + \delta_{in}n(t)} \left( (i - 1 + \delta_{in})x_{i-1}(t) - (i + \delta_{in})x_i(t) \right) + \alpha \mathbf{1}_{\{i=0\}} + \gamma \mathbf{1}_{\{i=1\}},$$

where we take  $x_{-1}(t) = 0$ , and write  $\mathbf{1}_{\mathcal{A}}$  for the indicator function which is 1 if the event  $\mathcal{A}$  holds and 0 otherwise.

Let  $i$  be fixed. We wish to take the expectation of both sides of (3.2). The only problem is with  $n(t)$  in the second term on the right hand side. For this, note that from a very weak form of (3.1), with probability  $1 - o(t^{-1})$  we have  $|n(t) - (\alpha + \gamma)t| = o(t^{3/5})$ . Now whatever value  $n(t)$  takes we have

$$\frac{\alpha + \beta}{t + \delta_{in}n(t)} (j + \delta_{in})x_j(t) = O(1)$$

for each  $j$ , so

$$\begin{aligned} \mathbf{E}\left(\frac{\alpha + \beta}{t + \delta_{in}n(t)} (j + \delta_{in})x_j(t)\right) &= \frac{\alpha + \beta}{t + \delta_{in}(\alpha + \gamma)t} (j + \delta_{in})\mathbf{E}x_j(t) + o(t^{-2/5}), \end{aligned}$$

and, taking the expectation of both sides of (3.2),

$$\begin{aligned} \mathbf{E}x_i(t+1) &= \mathbf{E}x_i(t) + \frac{\alpha + \beta}{t + \delta_{in}(\alpha + \gamma)t} \left( (i - 1 + \delta_{in})\mathbf{E}x_{i-1}(t) - (i + \delta_{in})\mathbf{E}x_i(t) \right) + \alpha \mathbf{1}_{\{i=0\}} + \gamma \mathbf{1}_{\{i=1\}} + o(t^{-2/5}). \end{aligned}$$

Let us write  $\bar{x}_i(t)$  for the ‘normalized expectation’  $\mathbf{E}x_i(t)/t$ . Recalling that  $c_1 = (\alpha + \beta)/(1 + \delta_{in}(\alpha + \gamma))$ , we have

$$(3.3) \quad \begin{aligned} (t+1)\bar{x}_i(t+1) - t\bar{x}_i(t) &= c_1 \left( (i - 1 + \delta_{in})\bar{x}_{i-1}(t) - (i + \delta_{in})\bar{x}_i(t) \right) + \alpha \mathbf{1}_{\{i=0\}} + \gamma \mathbf{1}_{\{i=1\}} + o(t^{-2/5}). \end{aligned}$$

Now let  $p_{-1} = 0$  and for  $i \geq 0$  define  $p_i$  by

$$(3.4) \quad \begin{aligned} p_i &= c_1 \left( (i - 1 + \delta_{in})p_{i-1} - (i + \delta_{in})p_i \right) + \alpha \mathbf{1}_{\{i=0\}} + \gamma \mathbf{1}_{\{i=1\}}. \end{aligned}$$

Our first claim is that for each  $i$  we have

$$(3.5) \quad \mathbf{E}(x_i(t)) = p_i t + o(t^{3/5})$$

as  $t \rightarrow \infty$ ; later we shall show that  $x_i(t)$  is concentrated around its mean, and then finally that the  $p_i$  follow the stated power law. To see (3.5), set  $\epsilon_i(t) = \bar{x}_i(t) - p_i$ . Then subtracting (3.4) from (3.3),

$$\begin{aligned} (t+1)\epsilon_i(t+1) - t\epsilon_i(t) &= c_1(i - 1 + \delta_{in})\epsilon_{i-1}(t) - c_1(i + \delta_{in})\epsilon_i(t) + o(t^{-2/5}), \end{aligned}$$

which we can rewrite as

$$(3.6) \quad \epsilon_i(t+1) = \frac{t - c_1(i + \delta_{in})}{t+1} \epsilon_i(t) + \Delta_i(t),$$

where  $\Delta_i(t) = c_1(i - 1 + \delta_{in})\epsilon_{i-1}(t)/(t+1) + o(t^{-7/5})$ .

To prove (3.5) we must show exactly that  $\epsilon_i(t) = o(t^{-2/5})$  for each  $i$ . We do this by induction on  $i$ ; suppose that  $i \geq 0$  and  $\epsilon_{i-1}(t) = o(t^{-2/5})$ , noting that  $\epsilon_{-1}(t) = 0$ , so the induction starts. Then  $\Delta_i(t) = o(t^{-7/5})$ , and from (3.6) one can check (for example by solving this equation explicitly for  $\epsilon_i(t)$  in terms of

$\Delta_i(t)$  that  $\epsilon_i(t) = o(t^{-2/5})$ . This completes the proof of (3.5).

Our next aim is to show that, with probability 1, we have

$$(3.7) \quad x_i(t)/t \rightarrow p_i,$$

as claimed in the statement of the theorem. To do this we show concentration of  $x_i(t)$  around its expectation using, as usual, the Azuma-Hoeffding inequality [4, 20] (see also [10]). This can be stated in the following form: if  $X$  is a random variable determined by a sequence of  $n$  choices, and changing one choice changes the value of  $X$  by at most  $\theta$ , then for any  $x \geq 0$  we have

$$(3.8) \quad \Pr(|X - \mathbf{E}X| \geq x) \leq 2e^{-x^2/(2n\theta^2)}.$$

To apply this let us first choose for each time step which operation (A), (B) or (C) to perform. Let  $\mathcal{A}$  be an event corresponding to one (infinite) sequence of such choices. Note that for almost all  $\mathcal{A}$  (in the technical sense of probability 1), the argument proving (3.5) actually gives

$$(3.9) \quad \mathbf{E}(x_i(t) | \mathcal{A}) = p_i t + o(t).$$

(We leave out the straightforward but somewhat technical details.)

Given  $\mathcal{A}$ , to determine  $G(t)$  it remains to choose at each step which old vertex (for (A) or (C)), or which old vertices (for (B)) are involved. There are at most  $2t$  old vertex choices to make. Changing one of these choices from  $v$  to  $v'$ , say, only affects the degrees of  $v$  and  $v'$  in the final graph. (To preserve proportional attachment at later stages we must redistribute later edges among  $v$  and  $v'$  suitably, but no other vertex is affected.) Thus  $x_i(t)$  changes by at most 2, and, applying (3.8), we have

$$\Pr\left(|x_i(t) - \mathbf{E}(x_i(t) | \mathcal{A})| \geq t^{3/4} | \mathcal{A}\right) \leq 2e^{-\sqrt{t}/16}.$$

Together with (3.9) this implies that (3.7) holds with probability one, proving the first part of the theorem. (Note that with a little more care we can probably replace (3.7) with  $x_i(t) = p_i t + O(t^{1/2} \log t)$ . Certainly our argument gives an error bound of this form in (3.5); the weaker bound stated resulted from replacing  $t^{1/2} \log t$  in (3.1) by  $o(t^{3/5})$  to simplify the equations. However, the technical details leading to (3.9) may become complicated if we aim for such a tight error bound.)

We now turn to the more substantial part of the result, determining the behaviour of the quantities  $p_i$  defined by (3.4).

Assuming  $\gamma < 1$ , we have  $\alpha + \beta > 0$  and hence  $c_1 > 0$ , so we can rewrite (3.4) as

$$(i + \delta_{in} + c_1^{-1})p_i = (i - 1 + \delta_{in})p_{i-1} + c_1^{-1}(\alpha \mathbf{1}_{\{i=0\}} + \gamma \mathbf{1}_{\{i=1\}}).$$

This gives  $p_0 = \alpha/(1 + c_1\delta_{in})$ ,

$$p_1 = (1 + \delta_{in} + c_1^{-1})^{-1} \left( \frac{\alpha\delta_{in}}{1 + c_1\delta_{in}} + \frac{\gamma}{c_1} \right),$$

and, for  $i \geq 1$ ,

$$(3.10) \quad \begin{aligned} p_i &= \frac{(i-1 + \delta_{in})_{i-1}}{(i + \delta_{in} + c_1^{-1})_{i-1}} p_1 \\ &= \frac{(i-1 + \delta_{in})!}{(i + \delta_{in} + c_1^{-1})!} \frac{(1 + \delta_{in} + c_1^{-1})!}{\delta_{in}!} p_1. \end{aligned}$$

Here, as usual, for  $x$  a real number and  $n$  an integer we write  $(x)_n$  for  $x(x-1)\cdots(x-n+1)$ . Also, we use  $x!$  for  $\Gamma(x+1)$  even if  $x$  is not an integer. We skip some detail in the derivations, as equations such as (3.4) clearly have unique solutions, and it is straightforward to check that the formulae we obtain do indeed give solutions. One can check that, as expected,  $\sum_{i=0}^{\infty} p_i = \alpha + \gamma$ ; there are  $(\alpha + \gamma + o(1))t$  vertices at large times  $t$ .

From (3.10) we see that as  $i \rightarrow \infty$  we have  $p_i \sim C_{IN} i^{-X_{IN}}$  with

$$X_{IN} = (\delta_{in} + c_1^{-1}) - (-1 + \delta_{in}) = 1 + 1/c_1,$$

as claimed.

For out-degrees the calculation is exactly the same after interchanging the roles of  $\alpha$  and  $\gamma$  and of  $\delta_{in}$  and  $\delta_{out}$ . Under this interchange  $c_1$  becomes  $c_2$ , so the exponent in the power law for out-degrees is  $X_{OUT} = 1 + 1/c_2$ , as claimed.

We now turn to more detailed results, considering in- and out-degree at the same time. Let  $n_{ij}(t)$  be the number of vertices of  $G(t)$  with in-degree  $i$  and out-degree  $j$ .

**THEOREM 3.2.** *Assume the conditions of Theorem 3.1 hold, that  $\alpha, \gamma < 1$ , and that  $\alpha\delta_{in} + \gamma\delta_{out} > 0$ . Let  $i, j \geq 0$  be fixed. Then there is a constant  $f_{ij}$  such that  $n_{ij}(t) = f_{ij}t + o(t)$  holds with probability 1. Furthermore, for  $j \geq 1$  fixed and  $i \rightarrow \infty$ ,*

$$(3.11) \quad f_{ij} \sim C_j i^{-X'_{IN}},$$

while for  $i \geq 1$  fixed and  $j \rightarrow \infty$ ,

$$(3.12) \quad f_{ij} \sim D_i j^{-X'_{OUT}},$$

where the  $C_j$  and  $D_i$  are positive constants,

$$X'_{IN} = 1 + 1/c_1 + c_2/c_1(\delta_{out} + \mathbf{1}_{\{\gamma\delta_{out}=0\}})$$

and

$$X'_{OUT} = 1 + 1/c_2 + c_1/c_2(\delta_{in} + \mathbf{1}_{\{\alpha\delta_{in}=0\}}).$$

Note that Theorem 3.2 makes statements about the limiting behaviour of the  $f_{ij}$  as one of  $i$  and  $j$  tends to infinity with the other fixed; there is no statement about the behaviour as  $i$  and  $j$  tend to infinity together in some way.

The proof of Theorem 3.2 follows the same lines as that of Theorem 3.1, but involves considerably more calculation, and is thus given as an appendix. The key difference is that instead of (3.10) we obtain a two dimensional recurrence relation (6.13) whose solution is much more complicated.

Before discussing the application of Theorems 3.1 and 3.2 to the web graph, note that if  $\delta_{out} = 0$  (as we shall assume when modelling the web graph), vertices born with out-degree 0 always have out-degree 0. Such vertices exist only if  $\gamma > 0$ . Thus  $\gamma\delta_{out} > 0$  is exactly the condition needed for the graph to contain vertices with non-zero out-degree which were born with out-degree 0. It turns out that when such vertices exist they dominate the behaviour of  $f_{ij}$  for  $j > 0$  fixed and  $i \rightarrow \infty$ . A similar comment applies to  $\alpha\delta_{in}$  with in- and out-degrees interchanged. If  $\alpha\delta_{in} = \gamma\delta_{out} = 0$  then every vertex not in  $G_0$  will have either in- or out-degree 0.

Note also for completeness that if  $\gamma\delta_{out} > 0$  then (3.11) holds for  $j = 0$  also. If  $\gamma = 0$  then  $f_{i0} = 0$  for all  $i$ . If  $\gamma > 0$  but  $\delta_{out} = 0$ , then among vertices with out-degree 0 (those born at a type (C) step), the evolution of in-degree is the same as among all vertices with non-zero out-degree taken together. It follows from Theorem 3.1 that in this case  $f_{i0} \sim C_0 i^{-X_{IN}}$ .

#### 4 Particular values

An interesting question is for which parameters (if any) our model reproduces the observed power laws for certain real-world graphs, in particular, the web graph.

For this section we take  $\delta_{out} = 0$  for the reasons explained in section 2. We assume that  $\alpha > 0$ , as otherwise there will only be finitely many vertices (the initial ones) with non-zero out-degree. As before, let  $c_1 = (\alpha + \beta)/(1 + \delta_{in}(\alpha + \gamma))$  and note that now  $c_2 = 1 - \alpha$ . We have shown that the power-law exponents are

$$X_{IN} = 1 + 1/c_1$$

for in-degree overall (or in-degree with out-degree fixed as 0),

$$X_{OUT} = 1 + 1/c_2$$

for out-degree overall, and that if  $\delta_{in} > 0$  we have exponents

$$X'_{IN} = 1 + 1/c_1 + c_2/c_1$$

for in-degree among vertices with fixed out-degree  $j \geq 1$ , and

$$X'_{OUT} = 1 + 1/c_2 + \delta_{in}c_1/c_2$$

for out-degree among vertices with fixed in-degree  $i \geq 0$ .

For the web graph, recently measured values of the first two exponents [13] are  $X_{IN} = 2.1$  and  $X_{OUT} = 2.7$ . (Earlier measurements in [3] and [23] gave the same value for  $X_{IN}$  but smaller values for  $X_{OUT}$ .) Our model gives these exponents if and only if  $c_2 = .59$ , so  $\alpha = .41$ , and  $c_1 = 1/1.1$ , so

$$\delta_{in} = \frac{1.1(\alpha + \beta) - 1}{1 - \beta}.$$

This equation gives a range of solutions: the extreme points are  $\delta_{in} = 0, \beta = .49, \gamma = .1$  and  $\delta_{in} = .24, \beta = .59, \gamma = 0$ .

As a test of the model one could measure the exponents  $X'_{IN}$  and  $X'_{OUT}$  (which may of course actually vary when the fixed out-/in-degree is varied). We obtain 2.75 for  $X'_{IN}$  and anything in the interval [2.7, 3.06] for  $X'_{OUT}$ .

#### 5 Other models

In many contexts, such as the web graph, it is clear that while preferential attachment is important, it is not the only, or perhaps even the main, reason for widely varying degrees. Another underlying cause which can produce this effect is the varying *fitness* or *attractiveness* of vertices or web pages; some web pages are just more interesting than others. This can be modelled mathematically using 'hidden variables'; each vertex has a random attractiveness, and preferential attachment depends on this and on degree. A model along these lines has been proposed by Bianconi and Barabási [8] (see also [15]).

Here we would like to propose a corresponding model for directed graphs: when a vertex  $v$  is created, two random numbers are associated with it,  $\lambda_v$  and  $\mu_v$ , its *in-* and *out-fitness*. Let us fix two distributions  $D_{IN}$  and  $D_{OUT}$  on the non-negative real numbers. (The simplest examples would be exponential or power-law distributions.) In what follows, for each new vertex  $v$  created we choose independently  $\lambda_v$  from  $D_{IN}$  and  $\mu_v$  from  $D_{OUT}$ , these choices being independent of all earlier choices. As before we fix  $\alpha, \beta, \gamma \geq 0$  with  $\alpha + \beta + \gamma = 1$ , and also  $\delta_{in}, \delta_{out} \geq 0$ . At time  $t_0 \geq 0$  we start with an initial graph  $G_0$  with  $t_0$  edges and  $n_0 \geq 1$  vertices, with certain fitnesses  $\lambda_v, \mu_v$  for the vertices  $v$  of  $G_0$ .

For  $t \geq t_0$  we form  $G(t+1)$  from  $G(t)$  as follows:

(A) With probability  $\alpha$ , add a new vertex  $v$  together

with an edge from  $v$  to an existing vertex  $w$ , where  $w$  is chosen according to  $\lambda(d_{in} + \delta_{in})$ .

(B) With probability  $\beta$ , add an edge from an existing vertex  $v$  to an existing vertex  $w$ , where  $v$  and  $w$  are chosen independently,  $v$  according to  $\mu(d_{out} + \delta_{out})$ , and  $w$  according to  $\lambda(d_{in} + \delta_{in})$ .

(C) With probability  $\gamma$ , add a new vertex  $w$  and an edge from an existing vertex  $v$  to  $w$ , where  $v$  is chosen according to  $\mu(d_{out} + \delta_{out})$ .

Here, to *choose  $v$  according to  $\lambda(d_{in} + \delta_{in})$*  means to choose  $v$  so that  $\Pr(v = v_i)$  is proportional to  $\lambda_{v_i}(d_{in}(v_i) + \delta_{in})$ , and to *choose  $v$  according to  $\mu(d_{out} + \delta_{out})$*  means to choose  $v$  so that  $\Pr(v = v_i)$  is proportional to  $\mu_{v_i}(d_{out}(v_i) + \delta_{out})$ , where the degrees are measured in  $G(t)$ .

Since the in- and out-degrees of vertices with different fitness will grow at different power-law rates, this model will produce some vertices of very high in-degree but low out-degree and vice-versa. This will be the topic of a forthcoming paper. Of course, one could also consider more general ‘preference functions’, depending on attractiveness and degree in a more complicated way, as well as a joint distribution for  $\lambda_v, \mu_v$ , combined with extra parameters as in [14]. However, there is always some benefit in keeping the model simple and the number of parameters small.

## References

- [1] W. Aiello, F. Chung and L. Lu, *A random graph model for power law graphs*, Experiment. Math., 10 (2001), pp. 53–66.
- [2] R. Albert and A. L. Barabási, *Statistical mechanics of complex networks*, arXiv:cond-mat/0106096 (2001)
- [3] R. Albert, H. Jeong and A. L. Barabási, *Diameter of the world-wide web*, Nature, 401 (1999), pp. 130–131.
- [4] K. Azuma, *Weighted sums of certain dependent variables*, Tôhoku Math. J., 3 (1967), pp. 357–367.
- [5] A.-L. Barabási and R. Albert, *Emergence of scaling in random networks*, Science, 286 (1999), pp. 509–512.
- [6] A.-L. Barabási, R. Albert and H. Jeong, *Mean-field theory for scale-free random networks*, Physica A, 272 (1999), pp. 173–187.
- [7] A.-L. Barabási, R. Albert and H. Jeong, *Scale-free characteristics of random networks: the topology of the world-wide web*, Physica A, 281 (2000), pp. 69–77.
- [8] G. Bianconi and A.-L. Barabási, *Competition and multiscaling in evolving networks*, cond-mat/0011029.
- [9] B. Bollobás, *Random Graphs, Second Edition*, Cambridge studies in advanced mathematics, vol. 73, Cambridge University Press, Cambridge, 2001, xvi+498pp.
- [10] B. Bollobás, *Martingales, isoperimetric inequalities and random graphs*, in *Combinatorics* (Eger, 1987), pp. 113–139, Colloq. Math. Soc. János Bolyai, 52, North-Holland, Amsterdam 1988.
- [11] B. Bollobás and O. M. Riordan, *The diameter of a scale-free random graph*, to appear in *Combinatorica*.
- [12] B. Bollobás, O. M. Riordan, J. Spencer, and G. Tusnády, *The degree sequence of a scale-free random graph process*, Random Structures and Algorithms, 18 (2001), pp. 279–290.
- [13] A. Broder, R. Kumar, F. Maghoul, P. Raghavan, S. Rajagopalan, R. Stata, A. Tomkins and J. Wiener, *Graph structure in the web*, Proc 9th WWW Conf. (2000), pp. 309–320.
- [14] C. Cooper and A. Frieze, *A general model of web graphs*, preprint.
- [15] S. N. Dorogovtsev and J. F. F. Mendes, *Evolution of random networks*, preprint.
- [16] P. Erdős and A. Rényi, *On random graphs. I*, Publ. Math. Debrecen, 6 (1959), pp. 290–297.
- [17] P. Erdős and A. Rényi, *On the evolution of random graphs*, Magyar Tud. Akad. Mat. Kutató Int. Közl., 5 (1960), pp. 17–61.
- [18] M. Faloutsos, P. Faloutsos and C. Faloutsos, *On power-law relationships of the internet topology*, SIGCOMM 1999, Comput. Commun. Rev., 29 (1999), 251.
- [19] E. N. Gilbert, *Random graphs*, Ann. Math. Statist., 30 (1959), pp. 1141–1144.
- [20] W. Hoeffding, *Probability inequalities for sums of bounded random variables*, J. Amer. Statist. Assoc., 58 (1963), pp. 13–30.
- [21] H. Jeong, B. Tombor, R. Albert, Z. N. Oltvai and A.-L. Barabási, *The large-scale organization of metabolic networks*, Nature, 407 (2000), pp. 651–654.
- [22] J. Kleinberg, R. Kumar, P. Raghavan, S. Rajagopalan and A. Tomkins, *The web as a graph: measurements, models, and methods*, COCOON 1999.
- [23] R. Kumar, P. Raghavan, S. Rajagopalan and A. Tomkins, *Extracting large scale knowledge bases from the web*, VLDB 1999.
- [24] R. Kumar, P. Raghavan, S. Rajagopalan, D. Sivakumar, A. Tomkins and E. Upfal, *Stochastic models for the web graph*, FOCS 2000.
- [25] M. E. J. Newman, *The structure of scientific collaboration networks*, Proc. Natl. Acad. Sci USA, 98 (2001), pp. 404–409.
- [26] M. E. J. Newman, S. H. Strogatz and D. J. Watts, *Random graphs with arbitrary degree distributions and their applications*, Phys. Rev. E, 64 (2001), 026118.
- [27] D. Osthus and G. Buckley, *Popularity based random graph models leading to a scale-free degree distribution*, preprint.
- [28] D. J. Watts and S. H. Strogatz, *Collective dynamics of ‘small-world’ networks*, Nature, 393 (1998), pp. 440–442.

## 6 Appendix: Proof of Theorem 3.2

In this appendix we give the proof of Theorem 3.2. Arguing as in the proof of Theorem 3.1 we see that for each  $i$  and  $j$  we have  $n_{ij}(t)/t \rightarrow f_{ij}$ , where the  $f_{ij}$

satisfy

$$(6.13) \quad \begin{aligned} f_{ij} &= c_1(i-1+\delta_{in})f_{i-1,j} - c_1(i+\delta_{in})f_{ij} \\ &+ c_2(j-1+\delta_{out})f_{i,j-1} - c_2(j+\delta_{out})f_{ij} \\ &+ \alpha \mathbf{1}_{\{i=0,j=1\}} + \gamma \mathbf{1}_{\{i=1,j=0\}}. \end{aligned}$$

Of course we take  $f_{ij}$  to be zero if  $i$  or  $j$  is  $-1$ . (At first sight there might seem to be a problem caused by the possibility that a vertex sends a loop to itself, increasing both its in- and out-degrees in one step. While this does complicate the equations for  $\mathbf{E}(n_{ij}(t))$ , it is easy to see that for fixed  $i$  and  $j$  the effect on this expectation is  $o(t)$ , so (6.13) holds exactly.)

We start by finding an expansion for  $f_{ij}$  when  $i \rightarrow \infty$  with  $j$  fixed.

The recurrence relation (6.13) is of the form

$$(6.14) \quad \mathcal{L}(\mathbf{f}) = \alpha \mathbf{1}_{\{i=0,j=1\}} + \gamma \mathbf{1}_{\{i=1,j=0\}}$$

for a linear operator  $\mathcal{L}$  on the two-dimensional array of coefficients  $\mathbf{f} = f_{ij}$ . We introduce the notation  $\mathcal{L}$  just as a convenience; to keep the formulae relatively simple, and reduce the number of cases, we wish to analyze the contributions to  $\mathbf{f}$  arising from the two terms on the right of (6.14) separately. The only property of  $\mathcal{L}$  needed to allow this separation is its linearity.

It is clear from the form of  $\mathcal{L}$  that there is a unique solution to equation (6.14). By linearity we can write

$$f_{ij} = g_{ij} + h_{ij}$$

where

$$(6.15) \quad \mathcal{L}(\mathbf{g}) = \alpha \mathbf{1}_{\{i=0,j=1\}}$$

and

$$(6.16) \quad \mathcal{L}(\mathbf{h}) = \gamma \mathbf{1}_{\{i=1,j=0\}}.$$

Let us first consider  $\mathbf{g}$ . As  $\alpha, \gamma < 1$  we have  $c_1, c_2 > 0$ , so setting

$$b_j = \delta_{in} + \frac{1}{c_1} + \frac{c_2}{c_1}(j + \delta_{out}),$$

dividing (6.15) through by  $c_1$  we obtain

$$(6.17) \quad \begin{aligned} (i+b_j)g_{ij} &= (i-1+\delta_{in})g_{i-1,j} \\ &+ \frac{c_2(j-1+\delta_{out})}{c_1}g_{i,j-1} + \frac{\alpha}{c_1} \mathbf{1}_{\{i=0,j=1\}}. \end{aligned}$$

Using (6.17), it is not hard to show that  $g_{ij} = 0$  for all  $i > 0$  if  $\alpha\delta_{in} = 0$ . For the moment, we shall therefore assume that  $\alpha\delta_{in} > 0$ .

Note that, from the boundary condition, we have  $g_{i0} = 0$  for all  $i$ . Thus for  $j = 1$  the second term on

the right of (6.17) disappears, and we see (skipping the details of the algebra) that

$$g_{i1} = a \frac{(i-1+\delta_{in})!}{(i+b_1)!}$$

where

$$a = \alpha \frac{(b_1-1)!}{c_1(\delta_{in}-1)!}$$

is a positive constant. (Here we have used  $\alpha\delta_{in} > 0$ .)

For  $j \geq 2$  the last term in (6.17) is always zero. Solving for  $g_{ij}$  by iteration, we get

$$(6.18) \quad g_{ij} = \frac{c_2(j-1+\delta_{out})}{c_1} \sum_{k=0}^i \frac{(i-1+\delta_{in})_{i-k}}{(i+b_j)_{i-k+1}} g_{k,j-1}.$$

Suppose that for some constants  $A_{jr}$  we have

$$(6.19) \quad g_{ij} = \sum_{r=1}^j A_{jr} \frac{(i-1+\delta_{in})!}{(i+b_r)!}$$

for all  $1 \leq j \leq j_0$  and all  $i \geq 0$ . Note that we have shown this for  $j_0 = 1$ , with  $A_{11} = a$ . Let  $j = j_0 + 1$ . Then, using (6.18) and (6.19), we see that

$$(6.20) \quad g_{ij} = \sum_{r=1}^{j-1} \frac{c_2(j-1+\delta_{out})}{c_1} A_{j-1,r} \sum_{k=0}^i \frac{(i-1+\delta_{in})!}{(i+b_j)_{i-k+1}(k+b_r)!}.$$

(The sums above are nested.) Now it is straightforward to verify that if  $0 < y < x$  and  $s$  is an integer with  $0 \leq s \leq i+1$ , then

$$(6.21) \quad \begin{aligned} &\sum_{k=s}^i \frac{1}{(i+x)_{i-k+1}(k+y)!} \\ &= \frac{1}{x-y} \left( \frac{1}{(i+y)!} - \frac{(s-1+x)!}{(i+x)!(s-1+y)!} \right). \end{aligned}$$

(For example one can use downwards induction on  $s$  starting from  $s = i+1$  where both sides are zero.) Combining (6.20) and the  $s = 0$  case of (6.21) we see that

$$\begin{aligned} g_{ij} &= \sum_{r=1}^{j-1} \frac{c_2(j-1+\delta_{out})}{c_1} A_{j-1,r} \frac{(i-1+\delta_{in})!}{b_j - b_r} \\ &\times \left( \frac{1}{(i+b_r)!} - \frac{(b_j-1)!}{(i+b_j)!(b_r-1)!} \right). \end{aligned}$$

(The last factor is of course inside the sum.) Collecting coefficients of  $1/(i + b_r)!$  for different values of  $r$ , and noting that  $b_j - b_r = (j - r)c_2/c_1$ , we see that (6.19) holds for  $j = j_0 + 1$ , provided that

$$A_{jr} = \frac{j - 1 + \delta_{out}}{j - r} A_{j-1,r}$$

for  $1 \leq r \leq j - 1$ , and

$$A_{jj} = - \sum_{r=1}^{j-1} \frac{j - 1 + \delta_{out}}{j - r} \frac{(b_j - 1)!}{(b_r - 1)!} A_{j-1,r}.$$

In fact we have the power law we are interested in (for  $\mathbf{g}$  rather than  $\mathbf{f}$ ) without calculating the  $A_{jr}$ . Observing only that  $A_{11} > 0$ , so  $A_{j1} > 0$  for every  $j \geq 1$ , the  $r = 1$  term dominates (6.19). Thus for any fixed  $j > 0$  we have

$$(6.22) \quad g_{ij} \sim C'_j i^{-1+\delta_{in}-b_1} = C'_j i^{-(1+1/c_1+c_2/c_1(1+\delta_{out}))}.$$

Having said that we do not need the  $A_{jr}$  for the power law, we include their calculation for completeness since it is straightforward. Skipping the rather unpleasant derivation, we claim that

$$A_{jr} = a(-1)^{r-1} \frac{(j - 1 + \delta_{out})!}{\delta_{out}!(j - 1)!} \binom{j - 1}{r - 1} \frac{(b_r - 1)!}{(b_1 - 1)!},$$

for the same constant  $a$  as above. This is easy to verify by induction on  $j$  using the relations above.

We now turn to  $\mathbf{h}$ , for which the calculation is similar. From (6.16) we have

$$(6.23) \quad (i + b_j)h_{ij} = (i - 1 + \delta_{in})h_{i-1,j} + \frac{c_2(j - 1 + \delta_{out})}{c_1} h_{i,j-1} + \frac{\gamma}{c_1} \mathbf{1}_{\{i=1, j=0\}}.$$

Again skipping much of the algebra, for  $j = 0$  we see that  $h_{00} = 0$ , while

$$h_{i0} = \gamma \frac{b_0!}{c_1 \delta_{in}!} \frac{(i - 1 + \delta_{in})!}{(i + b_0)!}$$

for all  $i \geq 1$ .

If  $\gamma \delta_{out} = 0$ , then  $h_{ij} = 0$  is zero for all  $j > 0$ , so let us now assume  $\gamma \delta_{out} > 0$ . This time the boundary condition implies that  $h_{0j} = 0$  for all  $j$ . For  $j \geq 1$  we thus have from (6.23) that

$$h_{ij} = \sum_{k=1}^i \frac{c_2(j - 1 + \delta_{out})}{c_1} h_{k,j-1} \frac{(i - 1 + \delta_{in})_{i-k}}{(i + b_j)_{i-k+1}}.$$

(The only difference from (6.18) is that the sum starts with  $k = 1$ .) Arguing as before, using the  $s = 1$  case of (6.21), we see that, for  $i \geq 1$  and  $j \geq 0$ ,

$$h_{ij} = \sum_{r=0}^j B_{jr} \frac{(i - 1 + \delta_{in})!}{(i + b_r)!},$$

where

$$B_{jr} = (-1)^r \gamma \frac{(j - 1 + \delta_{out})!}{j!(\delta_{out} - 1)!} \binom{j}{r} \frac{b_r!}{c_1 \delta_{in}!}.$$

(This makes sense as we are assuming that  $\delta_{out} > 0$ .) Here the  $r = 0$  term dominates, and we see that for each  $j \geq 0$  we have

$$(6.24) \quad h_{ij} \sim C''_j i^{-1+\delta_{in}-b_0} = C''_j i^{-(1+1/c_1+c_2\delta_{out}/c_1)}$$

as  $i \rightarrow \infty$ , for some positive constant  $C''_j$ . Returning now to  $\mathbf{f} = \mathbf{g} + \mathbf{h}$ , considering  $j \geq 1$  fixed and  $i \rightarrow \infty$  we see that when  $\gamma \delta_{out} > 0$ , the contribution from  $\mathbf{h}$  dominates, while if  $\gamma \delta_{out} = 0$ , this contribution is zero. Thus combining (6.22) and (6.24) proves (3.11).

The second part of Theorem 3.2, the proof of (3.12), follows by interchanging in- and out-degrees,  $\alpha$  and  $\gamma$  and  $\delta_{in}$  and  $\delta_{out}$ .